

Microwave background anisotropies from gravitational waves: the 1+3 covariant approach

Anthony Challinor[‡]

Astrophysics Group, Cavendish Laboratory, Madingley Road, Cambridge CB3 0HE, UK

Abstract. We present a 1+3 covariant discussion of the contribution of gravitational waves to the anisotropy of the cosmic microwave background radiation (CMB) in an almost-Friedmann-Robertson-Walker (FRW) universe. Our discussion is based in the covariant approach to perturbations in cosmology, which provides a physically transparent and gauge-invariant methodology for CMB physics. Applying this approach to linearised gravitational waves, we derive a closed set of covariant equations describing the evolution of the shear and the Weyl tensor, and the angular multipoles of the CMB intensity, valid for an arbitrary matter description and background spatial curvature. A significant feature of the present approach is that the normal mode expansion of the radiation distribution function, which arises naturally here, preserves the simple quadrupolar nature of the anisotropic part of the Thomson scattering source terms, and provides a direct characterisation of the power in the CMB at a given angular multipole, as in the recently introduced total angular momentum method. We provide the integral solution of the multipole equations, and analytic solutions for the shear and the Weyl tensor, for models with arbitrary spatial curvature. Numerical results for the CMB power spectrum in open models are also presented, which provide an independent verification of the calculations of other groups.

PACS numbers: 98.80-k, 98.70.Vc

1. Introduction

The cosmic microwave background radiation (CMB) plays an essential role in modern cosmology. Improved observations of the structure in the CMB should allow one to distinguish between rival theories of structure formation, and, in the case of the inflationary scenario, to determine the spectrum of primordial scalar (density) and tensor (gravitational wave) perturbations. Vector perturbations are expected to leave a significant imprint on the CMB only in the presence of active seeds, such as the defect models considered in [1].

The calculation of CMB anisotropies in specific cosmological models is now a large industry. For a representative sample of the literature dealing with the imprint of scalar perturbations, see e.g. [2]–[8]; for tensor perturbations see e.g. [9]–[19]. Early

[‡] A.D.Challinor@mrao.cam.ac.uk

calculational schemes were based on linear perturbation theory in the synchronous gauge [20], with Legendre expansions of the relativistic distribution functions [3, 5]. Some technical improvements in these schemes resulted from the use of gauge-invariant variables [6, 21], and, more recently, from improved normal mode expansions of the distribution functions [22, 23]. Adopting gauge-invariant variables frees one from the need to keep track of the implications of any residual gauge freedom in the calculation (the so called gauge problem; see e.g. [24, 25]), while improvements in the normal mode expansions have led to more direct characterisations of the anisotropy in vector and tensor modes, and a simpler form for the anisotropic Thomson scattering terms than is possible with the traditional Legendre expansion.

Recently, the 1+3 covariant and gauge-invariant approach to perturbations in cosmology, developed by Ellis and coworkers [24, 26] from the covariant hydrodynamics of Ehlers [27], has been applied to problems in CMB physics [28]–[36]. The 1+3 covariant approach provides a very convenient framework within which to study the exact intrinsic dynamics of cosmological models. The approach is formulated at the level of variables that are covariantly defined in the real universe, and are (exactly) gauge-invariant by construction, thus freeing one from many of the limitations of previous methodologies. Furthermore, the approach admits a covariant and gauge-invariant linearisation that allows linearised calculations to be performed in a very direct manner. In [32] a general 1+3 covariant scheme for the calculation of linearised CMB anisotropies was developed (see also [33, 36]). A complete, closed set of equations, that fully describe the linearised evolution of inhomogeneity and anisotropy in an almost-Friedmann-Robertson-Walker (FRW) universe with arbitrary spatial curvature, were presented. These equations were subsequently extended to include non-linear terms in [34], where some qualitative implications of non-linear effects were also noted. A significant feature of the covariant approach is the unified way in which scalar, vector and tensor perturbations are handled; the general perturbation equations are formulated without recourse to a decomposition into scalar, vector and tensor modes, although such a decomposition can still be made, if required, at a late stage in the calculation. In [32] the detailed equations describing scalar modes were derived from the unified equations, along with the necessary machinery to calculate the power spectrum of the CMB temperature anisotropy from a solution of these equations. It is our purpose here to report the extension of the 1+3 covariant analysis to tensor modes. We shall see that, in addition to the advantages of gauge-invariance and physical clarity that are evident in the scalar calculation, the tensor calculation highlights further advantages of the covariant approach over more traditional treatments. Namely, in the present approach a normal mode expansion of the radiation distribution function arises naturally that has the virtue of maintaining a direct characterisation of the power in the CMB at a given multipole ℓ , and of preserving the simple quadrupolar nature of that part of the Thomson scattering source terms that arises from the angular dependence of the Thomson cross section. Similar advantages are shared by the recently introduced total angular momentum method [22, 23].

The paper is arranged as follows. In section 2 we briefly review the 1+3 covariant approach to cosmological perturbations in an almost-FRW model, radiative transfer, and the decomposition of the CMB sky in angular multipoles. Section 3 begins with a discussion of the covariant description of gravitational waves in cosmology, and presents linearised wave equations, valid for a general matter stress-energy tensor, for the electric and magnetic parts of the Weyl tensor and the shear. These equations extend the perfect-fluid equations given in [37]. Following an expansion in tensor harmonics, in section 4 we give analytic solutions for the shear and the Weyl tensor during matter and radiation domination, extending the solutions given in [37] to the case of non-flat spatial sections. We also provide the integral solutions to the mode-expanded radiation multipole equations in the general case. In section 5 we derive the CMB power spectrum in terms of the mode-expanded multipole moments, and present numerical solutions in open CDM models, verifying the results in [18]. We close in section 6 with our conclusions. Appendix A contains useful new results on the symmetric trace-free representation of the tensor harmonics which we use in sections 4 and 5.

We employ standard general relativity with a $(+ - - -)$ metric signature. Our conventions for the Riemann and Ricci tensors are fixed by $2\nabla_{[a}\nabla_{b]}u_c = R_{abc}{}^d u_d$, and $R_{ab} \equiv R_{acb}{}^c$, where ∇_a denotes the covariant derivative. Round brackets denote symmetrisation on the enclosed indices, square brackets denote antisymmetrisation, and angle brackets denote the projected symmetric trace-free (PSTF) part. We use units with $c = G = 1$ throughout, so that the constant in the Einstein equation is $\kappa = 8\pi$. It should be noted that our convention for the metric signature, which coincides with that in our earlier work on CMB anisotropies in the 1+3 covariant approach [31, 32, 35], differs from much of the 1+3 covariant literature. It is straightforward to transform from our conventions to those in [34]: $g_{ab} \rightarrow -g_{ab}$, $\nabla_a \rightarrow \nabla_a$, and so $R_{abc}{}^d \rightarrow R_{abc}{}^d$. It follows that $R_{ab} \rightarrow R_{ab}$, but for the Ricci scalar $R \rightarrow -R$. The matter stress-energy tensor T_{ab} is unchanged, as is the fundamental velocity (vector) field u^a of the cosmological model. Raising or lowering an index incurs a sign change when transforming between signatures. The transformations of other key geometric objects are discussed in the text as they are introduced.

2. The 1+3 covariant approach to perturbations in cosmology

The essence of the 1+3 covariant approach is to identify a set of covariant variables which conveniently describe the inhomogeneity and anisotropy of the universe in a physically transparent (gauge-invariant) manner. Having chosen such variables, exact equations governing their dynamics can be found from the full field equations. Since the variables are chosen to have a direct physical interpretation, the physical content of their dynamical equations is often simple to extract. If required, the exact equations can be linearised around a chosen background to yield a simple set of perturbation equations that describe the evolution of physically relevant measures of the inhomogeneity and anisotropy of the universe in a very direct fashion.

The approach employs a 1+3 decomposition of geometric quantities with respect to a (fundamental) timelike velocity field u^a . We require that, in a general cosmological model, u^a be chosen in a physical manner such that in the FRW limit it correctly reduces to the preferred velocity. This restriction is necessary to ensure gauge-invariance of the approach. Tensors are decomposed irreducibly with respect to u^a into projected tensors which are orthogonal to u^a on every index. For example, the matter stress-energy tensor $T_{ab} = T_{(ab)}$ decomposes as§

$$T_{ab} = \rho u_a u_b + 2u_{(a} q_{b)} - p h_{ab} + \pi_{ab}, \quad (2.1)$$

which defines the energy density $\rho \equiv T_{ab} u^a u^b$, the energy flux $q_a \equiv h_a^b T_{bc} u^c$, the isotropic pressure $p \equiv -h^{ab} T_{ab}/3$, and the anisotropic stress $\pi_{ab} \equiv T_{\langle ab \rangle}$ which is the projected symmetric trace-free (PSTF) part of T_{ab} . Here, $h_{ab} \equiv g_{ab} - u_a u_b$ is the projection tensor into the instantaneous rest space of u^a (and g_{ab} is the metric of spacetime). In the FRW limit q_a and π_{ab} vanish since the stress-energy tensor is then restricted to perfect-fluid form.

It is convenient to introduce a projected covariant “derivative” D_a which acts on a general tensor $T_{b\dots}{}^{c\dots}$ according to

$$D_a T_{b\dots}{}^{c\dots} \equiv h_a^d h_b^e \dots h_f^c \dots \nabla_d T_{e\dots}{}^{f\dots}. \quad (2.2)$$

It should be noted that D_a is a derivation only when acting on projected objects. When u^a is irrotational (see below), D_a reduces to the covariant derivative in the hypersurfaces orthogonal to u^a . It is also convenient to introduce the derivative along the flow lines of u^a , denoted by an overdot, so that $\dot{T}_{b\dots}{}^{c\dots} = u^a \nabla_a T_{b\dots}{}^{c\dots}$. With these definitions, the covariant derivative of u^a decomposes as||

$$\nabla_a u_b = u_a A_b + \sigma_{ab} + \frac{1}{3} \Theta h_{ab} + \omega_{ab}, \quad (2.3)$$

which defines the acceleration $A_a \equiv \dot{u}_a$, the shear $\sigma_{ab} \equiv D_{\langle a} u_{b \rangle}$, the (volume) expansion scalar $\Theta = D^a u_a = 3H$, with H the local Hubble parameter, and the vorticity $\omega_{ab} \equiv D_{[a} u_{b]}$. Introducing a projected, totally antisymmetric tensor $\epsilon_{abc} \equiv \eta_{abcd} u^d$, where η_{abcd} is the totally antisymmetric tensor on spacetime, we can define the vorticity (projected) vector as the dual of ω_{ab} : $\omega_a \equiv \epsilon_{abc} \omega^{bc}/2$. It is also convenient to generalise the notion of the 3-dimensional curl to tensors $T_{ab\dots c}$ via

$$\text{curl } T_{ab\dots c} \equiv \epsilon_{de(a} D^d T_{b\dots c)}^e, \quad (2.4)$$

which is PSTF for $T_{ab\dots c}$ PSTF. The vorticity vector can then be written as $\omega_a = \text{curl } u_a/2$. The kinematic variables A_a , ω_a and σ_{ab} characterise anisotropy in the local expansion and thus vanish identically in an exact FRW universe. The projected gradient

§ To transform to the conventions of [34], one should make the replacements $q_a \rightarrow -q_a$, $\pi_{ab} \rightarrow \pi_{ab}$, as well as $u_a \rightarrow -u_a$ and $h_{ab} \rightarrow -h_{ab}$. Of course, ρ and p are unchanged.

|| To transform to the metric signature of [34], make the replacements $A_a \rightarrow -A_a$, $\sigma_{ab} \rightarrow -\sigma_{ab}$, $\omega_{ab} \rightarrow \omega_{ab}$, as well as $u_a \rightarrow -u_a$, $\nabla_a \rightarrow \nabla_a$ and $h_{ab} \rightarrow -h_{ab}$. It follows that for the vorticity vector $\omega_a \rightarrow \omega_a$. Our ω_a is physically minus that in [34] (rather than just any difference due to the metric signature), but coincides physically with that in [38].

of the expansion scalar, $D_a\Theta$, describes inhomogeneity in the expansion. Like the projected derivative of any covariant object, $D_a\Theta$ vanishes in the FRW limit.

The locally free part of the gravitational field is described by the Weyl tensor C_{abcd} . The ten degrees of freedom in C_{abcd} can be encoded in two PSTF tensors, E_{ab} and H_{ab} , denoted the electric and magnetic parts, respectively:[¶]

$$E_{ab} \equiv u^c u^d C_{acbd} \quad (2.5)$$

$$H_{ab} \equiv \frac{1}{2} \epsilon_{acd} C_{be}{}^{cd} u^e. \quad (2.6)$$

The electric part of the Weyl tensor plays an analogous role to the tidal tensor derived from the potential in Newtonian gravitation, while H_{ab} is essential for the propagation of gravitational waves. Indeed, the non-vanishing of the super-energy flux [39] $P_a \equiv \epsilon_{ac}{}^d E^{bc} H_{bd}$ for any timelike u^a forms the basis of Bel's first condition for the existence of gravitational radiation in an (exact) solution of the field equations (see e.g. [40]). Since the Weyl tensor vanishes in the FRW limit, E_{ab} and H_{ab} provide a covariant and gauge-invariant characterisation of the perturbation in the free gravitational field.

To construct a covariant linear perturbation theory about an FRW universe, those covariant variables that vanish in an exact FRW universe are regarded as being $O(\epsilon)$ in a smallness parameter ϵ . The linearisation procedure then consists of neglecting products of $O(\epsilon)$ variables in the dynamical equations. The exact dynamical equations, which follow from the field equations, are given in [34, 38]. Here, we shall only require the equations in linearised form around an FRW universe. There are seven propagation equations describing evolution along the flow lines of u^a :

$$\dot{\rho} = -(\rho + p)\Theta - D^a q_a \quad (2.7)$$

$$\dot{\Theta} = -\frac{1}{3}\Theta^2 - \frac{1}{2}\kappa(\rho + 3p) + D^a A_a \quad (2.8)$$

$$\dot{q}_a = -\frac{4}{3}\Theta q_a - (\rho + p)A_a + D_a p - D^b \pi_{ab} \quad (2.9)$$

$$\dot{\omega}_a = -\frac{2}{3}\Theta \omega_a + \frac{1}{2} \text{curl } A_a \quad (2.10)$$

$$\dot{\sigma}_{ab} = -\frac{2}{3}\Theta \sigma_{ab} - E_{ab} - \frac{1}{2}\kappa \pi_{ab} + D_{\langle a} A_{b\rangle} \quad (2.11)$$

$$\dot{E}_{ab} = -\Theta E_{ab} + \text{curl } H_{ab} + \frac{1}{2}\kappa \left[-(\rho + p)\sigma_{ab} - D_{\langle a} q_{b\rangle} + \dot{\pi}_{ab} + \frac{1}{3}\Theta \pi_{ab} \right] \quad (2.12)$$

$$\dot{H}_{ab} = -\Theta H_{ab} - \text{curl } E_{ab} - \frac{1}{2}\kappa \text{curl } \pi_{ab}, \quad (2.13)$$

and five constraint equations:

$$D^a \omega_a = 0 \quad (2.14)$$

$$D^a \sigma_{ab} - \text{curl } \omega_b - \frac{2}{3} D_b \Theta - \kappa q_b = 0 \quad (2.15)$$

$$D^a E_{ab} - \kappa \left(\frac{1}{3} \Theta q_b + \frac{1}{3} D_b \rho + \frac{1}{2} D^a \pi_{ab} \right) = 0 \quad (2.16)$$

[¶] Replace $E_{ab} \rightarrow -E_{ab}$ and $H_{ab} \rightarrow -H_{ab}$, and also $C_{acbd} \rightarrow -C_{acbd}$, $u^a \rightarrow u^a$, and $\epsilon_{acd} \rightarrow \epsilon_{acd}$ to transform to the conventions of [34].

$$D^a H_{ab} - \frac{1}{2} \kappa [2(\rho + p) \omega_b + \text{curl } q_b] = 0 \quad (2.17)$$

$$H_{ab} - \text{curl } \sigma_{ab} + D_{\langle a} \omega_{b \rangle} = 0, \quad (2.18)$$

which serve to constrain the initial data. It is straightforward to verify that the constraint equations are preserved by the propagation equations.

We describe the CMB radiation field via the specific intensity $I(E, e^a)$ which gives the intensity per unit solid angle along the (projected) direction e^a at photon energy E . The energy and propagation direction relative to u^a are related to the four-momentum of the photon by

$$p^a = E(e^a + u^a). \quad (2.19)$$

The photon distribution function $I(E, e^a)/E^3$ is frame-independent for given p^a . It is convenient to work with the energy-integrated intensity,

$$I(e^a) = \int_0^\infty I(E, e^a) dE, \quad (2.20)$$

since this determines the bolometric temperature along a given direction e^a . Denoting the fractional temperature difference from the all-sky average at some point by $\delta_T(e^a)$, we have

$$[1 + \delta_T(e^a)]^4 = \left(\frac{1}{4\pi} \int I(e^a) d\Omega \right)^{-1} I(e^a), \quad (2.21)$$

where the integration is over solid angles.

The anisotropy of the CMB intensity is conveniently described by the PSTF multipole moments $I_{A_\ell}^+$ in the covariant angular decomposition of $I(e^a)$:

$$I(e^a) = \sum_{\ell=0}^{\infty} \Delta_\ell^{-1} I_{A_\ell} e^{A_\ell}, \quad (2.22)$$

where e^{A_ℓ} is a convenient condensed notation for $e^{a_1} \dots e^{a_\ell}$, and we have defined

$$\Delta_\ell \equiv \frac{4\pi(-2)^\ell(\ell!)^2}{(2\ell+1)!}. \quad (2.23)$$

Including the factor of Δ_ℓ in (2.22) has the virtue of ensuring that the three lowest multipoles I , I_a and I_{ab} are simply the radiation energy density, energy flux, and anisotropic stress, respectively:

$$\rho^{(\gamma)} = I, \quad q_a^{(\gamma)} = I_a, \quad \pi_{ab}^{(\gamma)} = I_{ab}. \quad (2.24)$$

Propagation equations for the radiation multipoles follow from a multipole expansion of the Boltzmann equation [32, 34]. In linearised form, we have

$$\begin{aligned} \dot{I}_{A_\ell} + \frac{4}{3} \Theta I_{A_\ell} + D^b I_{bA_\ell} - \frac{\ell}{(2\ell+1)} D_{\langle a_\ell} I_{A_{\ell-1} \rangle} + \frac{4}{3} \delta_\ell^1 I_{A_a} - \frac{8}{15} \delta_\ell^2 I \sigma_{a_1 a_2} \\ = -n_e \sigma_T \left(I_{A_\ell} - \delta_\ell^0 I - \frac{4}{3} \delta_\ell^1 I v_{a_1}^{(b)} - \frac{1}{10} \delta_\ell^2 I_{a_1 a_2} \right), \end{aligned} \quad (2.25)$$

with $I_{A_\ell} = 0$ for $\ell < 0$. The projected vector $v_a^{(b)}$ which appears in the scattering term on the right of (2.25) is the relative velocity of the baryon frame, so that

⁺ We use a lumped index notation, $A_\ell = a_1 \dots a_\ell$.

$u_a^{(b)} = u_a + v_a^{(b)} + O(\epsilon^2)$ where $u_a^{(b)}$ is the baryon four-velocity. The electron number density in the baryon frame is denoted n_e , and σ_T is the Thomson cross section. We have ignored the small effect of polarization in (2.25). In [41] we give a full 1+3 covariant treatment of CMB polarization. There it is shown that the only effect of polarization on the intensity multipole equations is to replace $I_{a_1 a_2}$ by $I_{a_1 a_2} + 6\mathcal{E}_{a_1 a_2}$ in the final term on the right-hand side of (2.25), where $\mathcal{E}_{a_1 a_2}$ is a rank-2 PSTF tensor which describes the quadrupole moment of the electric polarization; see [23] for equivalent results in the total angular momentum approach. It is clear from the multipole equations that the complications due to the anisotropy of Thomson scattering affect only the $\ell = 2$ multipole equation. (This statement is also true of the further complications due the polarization dependence of Thomson scattering.) This simplicity, which is preserved by the normal-mode expansion described in section 3, ensures that the current methodology shares many of the advantages of the total angular momentum method over traditional Legendre techniques.

It follows from equation (2.21) that in linear theory, $\delta_T(e^a)$ has the multipole decomposition [28]

$$\delta_T(e^a) = \frac{\pi}{I} \sum_{\ell=1}^{\infty} \Delta_{\ell}^{-1} I_{A_{\ell}} e^{A_{\ell}}, \quad (2.26)$$

so that in a statistically isotropic ensemble the power spectrum of temperature anisotropies, C_{ℓ} , where

$$\langle \delta_T(e^a) \delta_T(e'^a) \rangle = \sum_{\ell=1}^{\infty} \frac{2\ell+1}{4\pi} C_{\ell} P_{\ell}(\cos \theta), \quad (2.27)$$

with $\cos \theta = -e^a e'_a$, can be determined from the covariance of the radiation multipoles [33]:

$$\left(\frac{\pi}{I}\right)^2 \langle I_{A_{\ell}} I^{B_{\ell'}} \rangle = \Delta_{\ell} C_{\ell} \delta_{\ell \ell'} h_{\langle A_{\ell}}^{\langle B_{\ell} \rangle}. \quad (2.28)$$

Here $h_{\langle A_{\ell}}^{\langle B_{\ell} \rangle} \equiv h_{\langle a_1}^{b_1} \dots h_{a_{\ell}}^{b_{\ell}} \rangle$, and large angle brackets denote an ensemble average.

3. Gravitational waves in an almost-FRW universe

The 1+3 covariant description of gravitational waves in a cosmological context has been considered by Hawking [42], and more recently by Dunsby, Bassett, & Ellis [37]. Linearised gravitational waves are described by the transverse degrees of freedom in the electric and magnetic parts of the Weyl tensor. Similarly, the shear and anisotropic stress are transverse so that

$$D^a E_{ab} = 0, \quad D^a H_{ab} = 0, \quad D^a \sigma_{ab} = 0, \quad D^a \pi_{ab} = 0 \quad (3.1)$$

at $O(\epsilon)$. Furthermore, the vorticity and all projected vectors that are generally $O(\epsilon)$ in an almost-FRW universe vanish at linear order. In particular, in a pure tensor mode the individual matter components all have the same four-velocity which defines the fundamental velocity u^a .

The covariant equations governing the propagation of linearised gravitational waves can be obtained from the equations of the previous section. The constraint equations (2.14–2.18) reduce to the single equation

$$H_{ab} = \text{curl } \sigma_{ab}. \quad (3.2)$$

which determines the magnetic part of the Weyl tensor from the shear. The propagation equations (2.7–2.13) reduce to

$$\dot{\rho} = -(\rho + p)\Theta \quad (3.3)$$

$$\dot{\Theta} = -\frac{1}{3}\Theta^2 - \frac{1}{2}\kappa(\rho + 3p) \quad (3.4)$$

$$\dot{\sigma}_{ab} = -\frac{2}{3}\Theta\sigma_{ab} - E_{ab} - \frac{1}{2}\kappa\pi_{ab} \quad (3.5)$$

$$\dot{E}_{ab} = -\Theta E_{ab} + \text{curl } H_{ab} + \frac{1}{2}\kappa \left[-(\rho + p)\sigma_{ab} + \dot{\pi}_{ab} + \frac{1}{3}\Theta\pi_{ab} \right] \quad (3.6)$$

$$\dot{H}_{ab} = -\Theta H_{ab} - \text{curl } E_{ab} - \frac{1}{2}\kappa \text{curl } \pi_{ab}, \quad (3.7)$$

Making use of the linearised identities [39], valid for an arbitrary first-order PSTF tensor S_{ab} ,

$$u^b \nabla_b (\text{curl } S_{ab}) = \text{curl } \dot{S}_{ab} - \frac{1}{3}\Theta \text{curl } S_{ab} \quad (3.8)$$

$$u^c \nabla_c (D^a S_{ab}) = D^a \dot{S}_{ab} - \frac{1}{3}\Theta D^a S_{ab} \quad (3.9)$$

$$D^a (\text{curl } S_{ab}) = \frac{1}{2} \text{curl } (D^a S_{ab}), \quad (3.10)$$

it is straightforward to verify that the constraints in equations (3.1) and (3.2) are consistent with the linearised propagation equations (3.3–3.7).

Inhomogeneous wave equations for the shear and the magnetic part of the Weyl tensor follow from differentiating equations (3.5) and (3.7) along the flow lines of u^a :

$$\ddot{\sigma}_{ab} + D^2 \sigma_{ab} + \frac{5}{3}\Theta \dot{\sigma}_{ab} + \left[\frac{1}{2}(4 - 3\gamma)\kappa\rho - \frac{K}{S^2} \right] \sigma_{ab} = -\kappa \left[\dot{\pi}_{ab} + \frac{2}{3}\Theta\pi_{ab} \right] \quad (3.11)$$

$$\ddot{H}_{ab} + D^2 H_{ab} + \frac{7}{3}\Theta \dot{H}_{ab} + 2 \left[(2 - \gamma)\kappa\rho - \frac{3K}{S^2} \right] H_{ab} = -\kappa \left[\text{curl } \dot{\pi}_{ab} + \frac{2}{3}\Theta \text{curl } \pi_{ab} \right], \quad (3.12)$$

where $D^2 \equiv D^a D_a$, and we have used the linearised identity

$$\text{curl } (\text{curl } S_{ab}) = D^2 S_{ab} + \frac{3K}{S^2} S_{ab} - \frac{3}{2} D_{\langle a} D^c S_{b \rangle c}. \quad (3.13)$$

Here, γ is defined in terms of the total energy density and pressure: $p = (\gamma - 1)\rho$, S is the scale factor of the universe [defined in a general almost-FRW model by $\dot{S} = HS$, and $D_a S = O(\epsilon)$], and $6K/S^2$ is the zero-order intrinsic curvature scalar on comoving hypersurfaces.

The solutions to the homogeneous wave equations in a $K = 0$ universe with constant γ were given in [37]. Solutions for $K \neq 0$ are given in section 4 in the radiation and matter eras. Solutions to the inhomogeneous equations can be obtained from those solutions by Green's method, but an accurate treatment of the matter-radiation

transition requires a numerical integration. The damping due to the anisotropic stress terms in the wave equations was discussed by Hawking for the simple viscous model $\pi_{ab} = \lambda \sigma_{ab}$, where λ is the viscosity [42]. During tight-coupling, the radiative contribution to π_{ab} for a general (long wavelength) perturbation is

$$I_{ab} = \frac{16}{27} I t_c (\sigma_{ab} + D_{\langle a} v_{b \rangle}^{(b)}), \quad (3.14)$$

where $t_c = (n_e \sigma_T)^{-1}$ is the photon mean free time. Including polarization changes the factor of 16/27 to 32/45 in this expression for I_{ab} , since the electric polarization quadrupole $\mathcal{E}_{ab} \approx I_{ab}/4$ during tight-coupling. The tensor $\sigma_{ab} + D_{\langle a} v_{b \rangle}^{(b)}$ is the shear in the baryon frame, so the effective (polarization-corrected) viscosity is $\lambda = 32 I t_c / 45$. Radiative viscous effects, as well as those due to other (collisionless) species such as neutrinos, lead to a small increase in the damping of gravitational waves over that due to redshift effects alone. However, unlike the case of scalar perturbations, the photon (and neutrino) quadrupole does not leave a significant imprint on the tensor CMB *temperature* power spectrum at high multipoles [19], since on the scales where damping due to anisotropic stresses could be significant, the tensor power spectrum is already insignificant due to sub-horizon redshifting of the gravity waves.

The wave equation for the electric part of the Weyl tensor is not closed (even for $\pi_{ab} = 0$) because of the presence of terms involving the shear:

$$\begin{aligned} \ddot{E}_{ab} + D^2 E_{ab} + \frac{7}{3} \Theta \dot{E}_{ab} + 2 \left[(2 - \gamma) \kappa \rho - \frac{3K}{S^2} \right] E_{ab} + \frac{1}{6} \kappa \rho \gamma \left[3 \frac{\dot{\gamma}}{\gamma} + (2 - 3\gamma) \Theta \right] \sigma_{ab} \\ = -\frac{1}{2} \kappa D^2 \pi_{ab} + \left[\frac{2}{3} \kappa \rho - \frac{3K}{S^2} \right] \kappa \pi_{ab} + \frac{5}{6} \Theta \kappa \dot{\pi}_{ab} + \frac{1}{2} \kappa \ddot{\pi}_{ab}. \end{aligned} \quad (3.15)$$

To eliminate the shear requires a further differentiation [42] (the resulting equation for $\pi_{ab} = 0$ was given in [37]). Although one cannot write down a closed wave equation for E_{ab} , the effective order of the equation is two since E_{ab} can be determined from the shear via equation (3.5), and the shear is determined by a wave equation.

3.1. Mode expansion in tensor harmonics

In linear perturbation theory it is convenient to expand the $O(\epsilon)$ variables in harmonic modes, since this converts the constraint equations into algebraic relations and the propagation equations into ordinary differential equations along the flow lines. For tensor perturbations, the appropriate harmonic functions can be derived from the tensor harmonics $Q_{ab}^{(k)}$. The zero-order properties of the tensor harmonics are discussed in Appendix A, where their explicit PSTF representation is derived for general K .

The electric and magnetic parts of the Weyl tensor, the shear, and the anisotropic stress can be expanded directly in the electric and magnetic parity tensor harmonics (see Appendix A):

$$E_{ab} = S^{-2} \sum_k k^2 (E_k Q_{ab}^{(k)} + \bar{E}_k \bar{Q}_{ab}^{(k)}) \quad (3.16)$$

$$H_{ab} = S^{-2} \sum_k k^2 (H_k Q_{ab}^{(k)} + \bar{H}_k \bar{Q}_{ab}^{(k)}) \quad (3.17)$$

$$\sigma_{ab} = S^{-1} \sum_k k (\sigma_k Q_{ab}^{(k)} + \bar{\sigma}_k \bar{Q}_{ab}^{(k)}) \quad (3.18)$$

$$\pi_{ab}^{(i)} = \rho^{(i)} \sum_k (\pi_k^{(i)} Q_{ab}^{(k)} + \bar{\pi}_k^{(i)} \bar{Q}_{ab}^{(k)}), \quad (3.19)$$

where the symbolic $\sum_k$ denotes a sum over the harmonic modes. For closed spatial sections ($K > 0$), the spectrum is discrete with $\nu \equiv [(k^2 + 3K)/|K|]^{1/2}$ an integer ≥ 3 . We use an overbar to distinguish the magnetic parity harmonics from the electric parity, and a subscript/superscript k to label (implicitly) the distinct eigenfunctions with degenerate eigenvalue k . For the representation in Appendix A, the label k represents the collection ν, ℓ, m , where ℓ and m describe the orbital angular momentum of the tensor harmonic. Functions of the eigenvalue alone will be denoted with k as an argument, for example, $A(k)$, or, equivalently, $A(\nu)$. The mode coefficients, such as E_k have $O(\epsilon^2)$ projected gradients by construction.

For the specific representation of the tensor harmonics constructed in Appendix A, the electric and magnetic parity tensor harmonics are related by a curl:

$$\text{curl } Q_{ab}^{(k)} = \frac{k}{S} \left(1 + \frac{3K}{k^2}\right)^{1/2} \bar{Q}_{ab}^{(k)}, \quad (3.20)$$

with an equivalent result for $\text{curl } \bar{Q}_{ab}^{(k)}$. From equation (3.2), we find that H_k is algebraic in $\bar{\sigma}_k$:

$$H_k = \left(1 + \frac{3K}{k^2}\right)^{1/2} \bar{\sigma}_k. \quad (3.21)$$

Note how the presence of curl terms leads to a coupling of the different polarization states. The constraint equation (3.2) [or, equivalently, (3.21)] allows one to eliminate H_{ab} from the discussion, leaving coupled first-order equations for the electric part of the Weyl tensor and the shear:

$$\frac{k^2}{S^2} \left(\dot{E}_k + \frac{1}{3} \Theta E_k \right) - \frac{k}{S} \left(\frac{k^2}{S^2} + \frac{3K}{S^2} - \frac{1}{2} \gamma \kappa \rho \right) \sigma_k = -\frac{1}{6} (3\gamma - 1) \kappa \rho \Theta \pi_k + \frac{1}{2} \kappa \rho \dot{\pi}_k \quad (3.22)$$

$$\frac{k}{S} \left(\dot{\sigma}_k + \frac{1}{3} \Theta \sigma_k \right) + \frac{k^2}{S^2} E_k = -\frac{1}{2} \kappa \rho \pi_k. \quad (3.23)$$

Combining these equations gives the harmonic expansion of the wave equation for the shear [equation (3.11)]:

$$\ddot{\sigma}_k + \Theta \dot{\sigma}_k + \left[\frac{k^2}{S^2} + \frac{2K}{S^2} - \frac{1}{3} (3\gamma - 2) \kappa \rho \right] \sigma_k = \kappa \rho \frac{S}{k} \left[\frac{1}{3} (3\gamma - 2) \Theta \pi_k - \dot{\pi}_k \right]. \quad (3.24)$$

For the radiation multipoles, I_{A_ℓ} , we expand in PSTF tensors $Q_{A_\ell}^{(k)}$ (and their overbarred counterparts) derived from the tensor harmonics via

$$Q_{A_\ell}^{(k)} = \left(\frac{S}{k} \right)^{\ell-2} D_{\langle a_1} \dots D_{a_{\ell-2}} Q_{a_{\ell-1} a_\ell}^{(k)}. \quad (3.25)$$

The factor of $(S/k)^{\ell-2}$ ensures that $\dot{Q}_{A_\ell}^{(k)} = O(\epsilon)$. For the ℓ -th multipole,

$$I_{A_\ell} = I \sum_k \beta_\ell^{-1} (I_k^{(\ell)} Q_{A_\ell}^{(k)} + \bar{I}_k^{(\ell)} \bar{Q}_{A_\ell}^{(k)}), \quad (3.26)$$

where, for later convenience, we have defined

$$\beta_\ell \equiv \prod_{n=2}^{\ell} \kappa_n, \quad (3.27)$$

$$\kappa_\ell \equiv [1 - (\ell^2 - 3)K/k^2]^{1/2}, \quad \ell \geq 2. \quad (3.28)$$

It follows that the photon anisotropic stress $\pi_k^{(\gamma)} = \kappa_2^{-1} I_k^{(2)}$. The $Q_{A_\ell}^{(k)}$ satisfy a number of differential identities (which are independent of any specific representation for the tensor harmonics), the most useful of which are

$$D^{a_\ell} Q_{A_\ell}^{(k)} = \frac{(\ell^2 - 4)}{\ell(2\ell - 1)} \frac{k}{S} \left[1 - (\ell^2 - 3) \frac{K}{k^2} \right] Q_{A_{\ell-1}}^{(k)} \quad (3.29)$$

$$D^2 Q_{A_\ell}^{(k)} = \frac{k^2}{S^2} \left[1 - (\ell - 2)(\ell + 3) \frac{K}{k^2} \right] Q_{A_\ell}^{(k)} \quad (3.30)$$

for $\ell \geq 2$, and the recursion relation

$$Q_{A_\ell}^{(k)} = \frac{k}{S} D_{\langle a_\ell} Q_{A_{\ell-1}}^{(k)} \quad (3.31)$$

for $\ell \geq 3$. In a closed universe, the $Q_{A_\ell}^{(k)}$ vanish for $\ell \geq \nu$, so that only modes with $\nu > \ell$ contribute to I_{A_ℓ} . The distinct processes of decomposing $I(e^a)$ into angular multipoles I_{A_ℓ} , followed by an expansion of the multipoles in derivatives of the tensor harmonics, combine to give a normal mode expansion of the radiation distribution function which may be regarded as the extension of the Legendre tensor approach, first introduced by Wilson [4], to non-scalar perturbations. In our opinion, it is one of the virtues of the 1+3 covariant approach that the physically significant decomposition into angular multipoles I_{A_ℓ} is kept distinct from the mathematically convenient harmonic mode decomposition. Maintaining this distinction allows one to describe the evolution of the anisotropy in a general cosmological model directly, without recourse to a harmonic splitting of the perturbations. Moreover, the present treatment is manifestly gauge-invariant, although we do not labour this point since gauge-invariance is less of an issue for tensor modes than for scalars.

Inserting the mode expansion, equation (3.26), into the propagation equation (2.25) for the multipoles I_{A_ℓ} , we find the mode-expanded multipole equations:

$$\begin{aligned} \dot{I}_k^{(\ell)} + \frac{k}{S} \left[\frac{(\ell + 3)(\ell - 1)}{(\ell + 1)(2\ell + 1)} \kappa_{\ell+1} I_k^{(\ell+1)} - \frac{\ell}{(2\ell + 1)} \kappa_\ell I_k^{(\ell-1)} \right] - \frac{8}{15} \frac{k}{S} \delta_\ell^2 \kappa_2 \sigma_k \\ = -n_e \sigma_T \left(I_k^{(\ell)} - \frac{1}{10} \delta_\ell^2 I_k^{(2)} \right), \end{aligned} \quad (3.32)$$

with similar equations for the barred variables. Massless neutrinos satisfy the same multipole equations but with the Thomson scattering terms omitted. If polarization is included, $I_k^{(2)}$ should be replaced by $I_k^{(2)} + 6\mathcal{E}_k^{(2)}$ in the last term on the right-hand side of (3.32), where $\mathcal{E}_k^{(2)}$ is the coefficient in the harmonic expansion of the electric polarization quadrupole; see [41] for further details. Equations (3.24) and (3.32), plus the massless neutrino equations, form a closed set of equations that can be solved to give

a complete description of the tensor contribution to the CMB anisotropy*. Analytic solutions for the shear and electric part of the Weyl tensor are given in the next section, assuming that anisotropic stress effects are negligible, as well as the integral solution for the radiation multipoles.

We end this section by noting that in kinetic theory, where the radiation multipoles may be non-vanishing for all ℓ , the familiar decomposition of the perturbations into scalar, vector and tensor modes is incomplete. A complete set of eigenfunctions for expanding a rank- ℓ PSTF tensor must include the equivalents of the scalar (rank-0), vector (rank-1), and tensor (rank-2) harmonics up to rank- ℓ . Since the perturbations in the matter stress-energy tensor, the kinematic variables, and the Weyl tensor are not affected by the higher-rank perturbations in linear theory (where perturbations of different rank evolve independently) the higher-rank perturbations in the distribution function cannot arise spontaneously. Any power that was initially in higher-rank perturbations of the photon distribution would thus be damped away due to Thomson scattering, leaving a negligible effect on the present day CMB anisotropy.

4. Analytic solutions for E_{ab} , σ_{ab} and $I_{A\ell}$

During tight-coupling, the photon anisotropic stress is suppressed by Thomson scattering, so the only significant contribution to π_{ab} is from the neutrinos. Assuming that this contribution is negligible, it is straightforward to find the analytic solutions for the Weyl tensor and the shear during radiation or matter domination. Anisotropic stresses can be included analytically by constructing the Green's functions from the solutions of the homogeneous equations given here.

It is convenient to use $x \equiv \sqrt{|K|}\eta$ as the independent variable, where η is conformal time. During radiation domination in an open universe, $HS = \sqrt{|K|} \coth x$, so that (3.24) becomes

$$\frac{d^2\sigma_k}{dx^2} + 2 \coth x \frac{d\sigma_k}{dx} + \left(\nu^2 + 1 - \frac{2}{\sinh^2 x} \right) \sigma_k = 0. \quad (4.1)$$

The solution of this equation which is regular as $\eta \rightarrow 0$ is

$$\sigma_k \propto \frac{\cos \nu x}{\sinh x} - \frac{\sin \nu x \coth x}{\nu \sinh x}, \quad (4.2)$$

and the irregular solution is

$$\sigma_k \propto \frac{\sin \nu x}{\sinh x} + \frac{\cos \nu x \coth x}{\nu \sinh x}. \quad (4.3)$$

These solutions for the shear determine the electric part of the Weyl tensor from equation (3.23). The regular solution evaluates to

$$E_k \propto \frac{1}{\sqrt{\nu^2 + 3} \sinh x} \left[\frac{\sin \nu x}{\nu} (\nu^2 + 1 - \coth^2 x) + \cos \nu x \coth x \right], \quad (4.4)$$

* When polarization is included, additional multipole hierarchies must be included for the electric and magnetic multipoles of the polarization to close the equations [41].

and the irregular solution to

$$E_k \propto \frac{-1}{\sqrt{\nu^2 + 3} \sinh x} \left[\frac{\cos \nu x}{\nu} (\nu^2 + 1 - \coth^2 x) - \sin \nu x \coth x \right]. \quad (4.5)$$

For $k\eta \ll 1$ (which implies $\nu x \ll 1$ and $x \ll 1$), the mode is well outside the horizon since $k\eta \ll 1$ implies that $HS/k \approx 2/(k\eta)$. When this condition is met the electric part of the Weyl tensor is approximately constant, with equation (4.4) giving $E_k \propto 2(\nu^2 + 1)/[3\sqrt{(\nu^2 + 3)}]$. The solutions for a closed universe can be obtained from those in an open universe with the substitutions $x \rightarrow ix$ and $\nu \rightarrow -i\nu$.

During matter domination in an open universe, $HS = \sqrt{|K|} \coth(x/2)$, so that (3.24) becomes

$$\frac{d^2 \sigma_k}{dx^2} + 2 \coth(x/2) \frac{d\sigma_k}{dx} + \left(\nu^2 + 1 - \frac{1}{\sinh^2(x/2)} \right) \sigma_k = 0. \quad (4.6)$$

The regular solution of this equation is

$$\sigma_k \propto \frac{1}{\sinh^2(x/2)} \left\{ \frac{\sin \nu x}{2\nu} [4\nu^2 + 1 - 3 \coth^2(x/2)] + 3 \cos \nu x \coth(x/2) \right\}, \quad (4.7)$$

and the irregular solution is

$$\sigma_k \propto \frac{1}{\sinh^2(x/2)} \left\{ \frac{\cos \nu x}{2\nu} [4\nu^2 + 1 - 3 \coth^2(x/2)] - 3 \sin \nu x \coth(x/2) \right\}. \quad (4.8)$$

Using equation (3.23), the regular solution for the electric part of the Weyl tensor during matter domination is

$$E_k \propto \frac{1}{\sqrt{\nu^2 + 3} \sinh^2(x/2)} \left\{ \frac{3 \sin \nu x}{2\nu} \coth(x/2) [2\nu^2 + 1 - \coth^2(x/2)] - \cos \nu x [2\nu^2 + 2 - 3 \coth^2(x/2)] \right\}, \quad (4.9)$$

and the irregular solution is

$$E_k \propto \frac{1}{\sqrt{\nu^2 + 3} \sinh^2(x/2)} \left\{ \frac{3 \cos \nu x}{2\nu} \coth(x/2) [2\nu^2 + 1 - \coth^2(x/2)] + \sin \nu x [2\nu^2 + 2 - 3 \coth^2(x/2)] \right\}. \quad (4.10)$$

On superhorizon scales, $E_k \propto 2(\nu^2 + 1)(4\nu^2 + 1)/[5\sqrt{(\nu^2 + 3)}]$ is approximately constant in the regular mode. In both the matter and radiation eras, $\sigma_k = O(x)$ in the regular modes on superhorizon scales. It follows from (3.21) that on superhorizon scales the Weyl tensor is dominated by the contribution from its electric part. The solutions given here generalise the $K = 0$ solutions derived in [37]; in the limit $K \rightarrow 0$ we reproduce their results.

We can provide a formal solution to equation (3.32) for the intensity multipoles by noting that the homogeneous equations (obtained by setting $n_e = 0$ and $\sigma_k = 0$) are solved by the functions $\ell(\ell - 1)\Phi_\ell^\nu(x)/\sinh^2(x)$ in an open universe, where $x \equiv \sqrt{|K|}(\eta_R - \eta)$ with η_R the conformal time at our current position R . Here, $\Phi_\ell^\nu(x)$ are

the ultra-spherical Bessel functions [6]; see Appendix A also. Defining the optical depth back to cosmic time t , with $t = t_R$ at R :

$$\tau(t) \equiv \int_t^{t_R} n_e \sigma_T dt', \quad (4.11)$$

the integral solution for $I_k^{(\ell)}$ in an open universe is

$$I_k^{(\ell)} = \frac{4\ell(\ell-1)}{[(\nu^2+1)(\nu^2+3)]^{1/2}} \int^{t_R} dt' e^{-\tau} \left(\frac{k}{S} \sigma_k + \frac{3}{16} n_e \sigma_T \pi_k^{(\gamma)} \right) \frac{\Phi_\ell^\nu(x)}{\sinh^2 x}. \quad (4.12)$$

For a closed universe, the hyperbolic functions should be replaced by their trigonometric counterparts and $\nu^2 + n$ by $\nu^2 - n$, where n is an integer. Equation (4.12) can be shown to be equivalent to the tensor integral solution derived in [23] with the total angular momentum method. The shear term in (4.12) arises from the cumulative redshifting effect of shear along the line of sight. The contribution to the anisotropy at R from the redshift incurred for a given increment along the null geodesic is weighted by the probability $e^{-\tau}$ that a photon subsequently does not scatter before arriving at R . The term involving the radiation anisotropic stress (intensity quadrupole) in equation (4.12) describes the cumulative effect of the anisotropic scattering into the beam along the line of sight. (Inclusion of polarization gives an additional contribution to this term so that $\pi_k^{(\gamma)}$ should be replaced by $\pi_k^{(\gamma)} + 6\kappa_2^{-1} \mathcal{E}_k^{(2)}$. This term has only a very small effect on the intensity multipoles over the range of ℓ where the magnitudes of the I_{A_ℓ} are appreciable.) The geometric factor $\Phi_\ell^\nu(x)/\sinh^2 x$ arises from the projection of the contraction $Q_{ab}^{(k)} e^a e^b$ of the tensor harmonic with the projected photon direction at comoving distance $x/\sqrt{|K|}$ back along the line of sight. This follows quite generally from noting that, for any representation of the tensor harmonics, $Q_{ab}^{(k)} e^a e^b$ at x down the line of sight is given in terms of the $Q_{A_\ell}^{(k)} e^{A_\ell}$ at the observation point by

$$Q_{ab}^{(k)} e^a e^b|_x = \frac{4\pi}{[(\nu^2+1)(\nu^2+3)]^{1/2}} \sum_{\ell=2}^{\infty} \frac{\ell(\ell-1)}{\Delta_\ell \beta_\ell} \frac{\Phi_\ell^\nu(x)}{\sinh^2 x} Q_{A_\ell}^{(k)} e^{A_\ell}|_R. \quad (4.13)$$

This result, which can be regarded as the PSTF extension of the Rayleigh plane wave result to eigentensors of the Laplacian on a manifold of constant curvature, can be verified by comparing the derivatives of $Q_{ab}^{(k)} e^a e^b$ at $x = 0$ with those computed from the following recursion relation for the $Q_{A_\ell}^{(k)} e^{A_\ell}$:

$$e^b D_b (Q_{A_\ell}^{(k)} e^{A_\ell}) = \frac{k}{S} \left[Q_{A_{\ell+1}}^{(k)} e^{A_{\ell+1}} - \frac{(\ell^2 - 4)}{(4\ell^2 - 1)} \kappa_\ell^2 Q_{A_{\ell-1}}^{(k)} e^{A_{\ell-1}} \right]. \quad (4.14)$$

In a closed universe the sum over ℓ in equation (4.13) is restricted to the range $2 \leq \ell \leq \nu - 1$.

5. The CMB power spectrum

To determine the CMB power spectrum from the $I_\ell^{(k)}$ we require the $Q_{A_\ell}^{(k)}$ at the observation point R . For this purpose, it is convenient to consider the variation of $Q_{ab}^{(k)} e^a e^b$ down the line of sight for the chosen representation of the harmonics, and to

compare this result with (4.13) which is independent of the specific representation. For the representation of the tensor harmonics given in Appendix A, we find for the electric parity harmonics

$$Q_{ab}^{(k)} e^a e^b|_x = \frac{1}{\nu \sqrt{2(\nu^2 + 1)}} \left[\frac{(\ell + 2)!}{(\ell - 2)!} \right]^{1/2} \frac{\Phi_\ell^\nu(x)}{\sinh^2 x} \mathcal{Q}_{A_\ell} e^{A_\ell}|_R, \quad (5.1)$$

in an open universe, and $\bar{Q}_{ab}^{(k)} e^a e^b|_x = 0$ for the magnetic parity harmonics. Here, $\mathcal{Q}_{A_\ell}$ is a rank- ℓ PSTF tensor encoding the $2\ell + 1$ degrees of freedom in the tensor harmonic with ℓ units of orbital angular momentum; see the discussion in Appendix A.3. Comparing equations (5.1) and (4.13), and using the orthogonality of the $e^{\langle A_\ell \rangle}$, we find

$$Q_{A_{\ell'}}^{(k)}|_R = \frac{1}{4\pi} \left[\frac{(\nu^2 + 3)}{2\nu^2} \right]^{1/2} \left[\frac{(\ell + 2)(\ell + 1)}{\ell(\ell - 1)} \right]^{1/2} \Delta_{\ell\beta_\ell} \mathcal{Q}_{A_\ell} \delta_{\ell\ell'}, \quad (5.2)$$

for the electric parity tensor harmonics, and $\bar{Q}_{A_{\ell'}}^{(k)}|_R = 0$ for magnetic parity. Although the magnetic parity harmonics do not contribute to the radiation anisotropy at R , they do contribute on other integral curves of u^a . The contribution of the electric parity harmonics undergoes a compensating change to ensure statistical homogeneity.

Following standard practice we assume that on the scales of cosmological interest today there is negligible power in the irregular modes given in section 4 during the radiation era. This assumption is justified by the highly squeezed quantum state predicted at the end of inflation [43, 44]. Since E_k is constant outside the horizon for the regular solution, it is convenient to parameterise the radiation multipoles in terms of this constant:

$$I_k^{(\ell)} = \frac{(\nu^2 + 3)}{(\nu^2 + 1)} T^{(\ell)}(\nu) \lim_{k\eta \rightarrow 0} E_k, \quad (5.3)$$

in an open universe. $T^{(\ell)}(\nu)$ is the radiation transfer function, and the ν -dependent factor is chosen to conform with treatments based on the metric perturbation variable (e.g. [23]). The primordial tensor power spectrum $\mathcal{P}(\nu)$ is defined in terms of the covariance matrix for the E_k in the limit $k\eta \rightarrow 0$:

$$\langle E_k E_{k'} \rangle = \langle \bar{E}_k \bar{E}_{k'} \rangle \propto \frac{1}{|K|^{3/2}} \frac{(\nu^2 + 1)^2}{(\nu^2 + 3)^2} \frac{\mathcal{P}(\nu)}{\nu(\nu^2 + 1)} \delta_{kk'}, \quad (5.4)$$

where $\delta_{kk'}$ is the symbolic delta function appropriate to the sum over tensor harmonics; see Appendix A. In the $K \rightarrow 0$ limit, $\mathcal{P}(\nu) \rightarrow k^{n_T}$ for a tensor spectral index n_T . Making use of equations (2.28) and (5.2), we find that the CMB power spectrum evaluates to

$$C_\ell = \frac{1}{16} \frac{(\ell + 2)(\ell + 1)}{\ell(\ell - 1)} \int_0^\infty \frac{\nu d\nu}{(\nu^2 + 1)} \frac{(\nu^2 + 3)}{2\nu^2} \mathcal{P}(\nu) [T^{(\ell)}(\nu)]^2 \quad (5.5)$$

in an open universe. For closed models the integral over ν should be replaced by a discrete sum over integral $\nu \geq \ell + 1$. Note that the CMB power spectrum at multipole ℓ depends only on the transfer functions at the same multipole; a feature not shared by schemes based on simple Legendre expansions of the distribution functions (e.g. [14]).

With these conventions, the contribution to the magnitude (squared) of the electric part of the Weyl tensor from superhorizon modes is

$$\left(\frac{S}{k}\right)^4 \frac{\partial}{\partial \ln k} \langle E_{ab} E^{ab} \rangle = \frac{1}{2\pi} \frac{(\nu^2 + 4)(\nu^2 + 1)}{\nu^2(\nu^2 + 3)} \mathcal{P}(\nu). \quad (5.6)$$

For $\nu \rightarrow 0$ in open models, the limit $\mathcal{P}(\nu) \rightarrow 0$ is required to avoid an infrared divergence in the C_ℓ and the electric part of the Weyl tensor. (In bubble models, $\mathcal{P}(\nu) \rightarrow \nu$ for $\nu \ll 1$ [45].)

The C_ℓ due to gravitational waves in the CDM model are plotted in Figure 1 for three values of the current-day matter density: $\Omega_0 = 0.1, 0.4$ and 1 . The Hubble constant $h = 0.7$ ($H_0 = 100h \text{ kms}^{-1}\text{Mpc}^{-1}$), the baryon fraction $\Omega_b h^2 = 0.0125$, and there is no reionisation or cosmological constant. The primordial power spectrum is the minimal scale-invariant spectrum $\mathcal{P}(\nu) = \tanh(\pi\nu/2)$ [45]. For the flat model ($\Omega_0 = 1$), this $\mathcal{P}(\nu) \rightarrow 1$ corresponding to $n_T = 0$. The power spectra were calculated by modifying CMBFAST [8] to solve the 1+3 covariant equations given in this paper. The results are in excellent agreement with those given in [18] which employed the total angular momentum method. Notable features in Figure 1, which are discussed further in [18], include the angular scaling brought about by changes in Ω_0 (variations in the angular diameter distance), the curvature cutoff in the large scale CMB power spectrum in low density universes (the curve for $\Omega_0 = 0.1$), and the enhancement in power on large scales, compared to the flat case, for higher values of the density parameter (the curve for $\Omega_0 = 0.4$), due to the growth of the shear in the longest wavelength modes once they enter the horizon.

6. Conclusion

We have developed the 1+3 covariant results necessary to calculate the temperature anisotropy of the CMB due to gravitational waves in general almost-FRW models. The covariant and gauge-invariant approach is well suited to describing gravitational waves in a cosmological context, since the emphasis is placed on the physically relevant Weyl tensor and shear, rather than the perturbation in the metric tensor. In the covariant calculation the intensity anisotropy is decomposed into irreducible form before linearisation around an FRW background, so the approach is also well suited to studying non-linear radiative effects (see [34]). For the linear calculation, the expansion of the radiation intensity in PSTF tensor-valued multipoles ensures that the subsequent mode expansions of the radiation variables for scalar, vector and tensor modes naturally generalise the harmonic expansions of the rank-0, 1 and 2 tensors which describe inhomogeneity and anisotropy in the geometry. The same is true for the description of the polarization of the CMB, as we show elsewhere [35, 41]. The technique should also allow a straightforward derivation of the multipole hierarchies for massive particles in difficult cases, such as non-flat models with vector or tensor perturbations.




Figure 1. Intensity multipoles from gravitational waves in CDM models with $h = 0.7$, $\Omega_b h^2 = 0.0125$, and no reionisation or cosmological constant. The C_ℓ are shown for two open models, $\Omega_0 = 0.1$ and 0.4 , and the flat model. The primordial power spectrum is $\mathcal{P}(\nu) = \tanh(\pi\nu/2)$. The curves are normalised at C_{10} .

Acknowledgments

The author acknowledges Queens' College, Cambridge, for support in the form of a Research Fellowship, and valuable help from Antony Lewis who made the modifications to the CMBFAST code.

Appendix A. The tensor harmonics

The tensor harmonics $Q_{ab}^{(k)}$ are zero-order, second-rank PSTF tensors satisfying

$$D^2 Q_{ab}^{(k)} = \frac{k^2}{S^2} Q_{ab}^{(k)}, \quad (\text{A.1})$$

where the equality holds at zero-order. They are constructed to satisfy the zero-order relations

$$D^a Q_{ab}^{(k)} = 0, \quad \dot{Q}_{ab}^{(k)} = 0, \quad (\text{A.2})$$

which are consistent with (A.1) at zero-order. With the definition $\nu \equiv [(k^2 + 3K)/|K|]^{1/2}$, the spectrum of regular, normalisable solutions for open and flat background models has $\nu \geq 0$, while in closed models ν takes integral values ≥ 3 . Supercurvature modes in open models (see e.g. [46]) correspond to $0 > \nu^2 \geq -3$. For $\nu^2 \gg 1$ the mode is well inside the curvature radius and k is an effective comoving wavenumber. We label the tensor harmonics with a lumped superscript (k) which should be understood to distinguish degenerate solutions of (A.1).

In the text we make use of a specific PSTF representation of the tensor harmonics. For this representation it is convenient to “coordinatise” the spacetime as follows. Starting at some arbitrary point R (which can conveniently be taken to be our current location), we construct spatial geodesics with tangent vectors $e^a(\chi)$, the generators of which at R are the set of projected unit vectors $\{e^a|_R\}$ at R :

$$e^a D_a e^b = 0, \quad u^a e_a = 0, \quad \frac{dx^a}{d\chi} = -S e^a, \quad (\text{A.3})$$

with $\chi \geq 0$ (the equality holding at R). In the absence of vorticity the curves $x^a(\chi)$ lie in the flow-orthogonal hypersurface through R . A projected vector field e^a and a scalar field χ can then be constructed in the neighbourhood of R from their restriction on the hypersurface defined by $\{e^a|_R, \chi\}$ via

$$\dot{e}^{(a)} = 0, \quad \dot{\chi} = 0. \quad (\text{A.4})$$

The fields e^a and χ satisfy equations (A.3) at zero-order in an almost-FRW universe. To the same order, the restriction of e^a to the past lightcone through R is the projected direction of a photon that passes through R , and χ is the zero-order conformal lookback time.

Appendix A.1. Electric parity harmonics

For the electric parity tensor harmonics we start with the most general electric parity, rank-two PSTF tensor [47] which is constant along the flow lines of u^a :

$$Q_{ab}^{(k)} = T_1(x) \left(e_a e_b \mathcal{Q}_{C_\ell} e^{C_\ell} + \frac{1}{2} \mathcal{H}_{ab} \mathcal{Q}_{C_\ell} e^{C_\ell} \right) + T_2(x) e_{(a} \mathcal{H}_{b)}^{C_\ell} \mathcal{Q}_{C_\ell} e^{C_{\ell-1}} + T_3(x) [\mathcal{Q}_{ab C_{\ell-2}} e^{C_{\ell-2}}]^{\text{TT}}, \quad \ell \geq 2. \quad (\text{A.5})$$

Here, $\mathcal{Q}_{A_\ell}$ is a rank- ℓ PSTF tensor field satisfying the zero-order equations

$$e^b D_b \mathcal{Q}_{A_\ell} = 0, \quad \dot{\mathcal{Q}}_{A_\ell} = 0, \quad (\text{A.6})$$

which determine the tensor field $\mathcal{Q}_{A_\ell}$ from its value at the point R , and $T_i(x)$, $i = 1, 2, 3$, are scalar functions of $x \equiv \sqrt{|K|}\chi$. The (screen) projection tensor $\mathcal{H}_{ab}$ is

$$\mathcal{H}_{ab} \equiv h_{ab} + e_a e_b, \quad (\text{A.7})$$

and the notation $[A_{ab}]^{\text{TT}}$ denotes the transverse traceless part of the second-rank tensor A_{ab} :

$$[A_{ab}]^{\text{TT}} \equiv \mathcal{H}_a^{c_1} \mathcal{H}_b^{c_2} A_{c_1 c_2} - \frac{1}{2} \mathcal{H}_{ab} \mathcal{H}^{c_1 c_2} A_{c_1 c_2}. \quad (\text{A.8})$$

Demanding that $D^a Q_{ab}^{(k)} = 0$ determines T_2 and T_3 in terms of T_1 . In an open universe we find

$$T_2(x) = \frac{-2}{(\ell+1) \sinh^2 x} \frac{d}{dx} [\sinh^3 x T_1(x)], \quad (\text{A.9})$$

$$T_3(x) = \frac{-\ell}{(\ell+2)} T_1(x) - \frac{1}{(\ell+2) \sinh^2 x} \frac{d}{dx} [\sinh^3 x T_2(x)]. \quad (\text{A.10})$$

Equation (A.1) gives three further equations, which are satisfied by virtue of equations (A.9) and (A.10) provided that

$$\frac{d^2}{dx^2}T_1(x) + 6 \coth x \frac{d}{dx}T_1(x) + \left[\nu^2 + 9 - \frac{(\ell^2 + \ell - 6)}{\sinh^2 x} \right] T_1(x) = 0. \quad (\text{A.11})$$

The regular, normalisable solution of this equation is

$$T_1(x) = N(\nu) \frac{\Phi_\ell^\nu(x)}{\sinh^2 x}, \quad (\text{A.12})$$

with $\nu \geq 0$ (subcurvature). The $\Phi_\ell^\nu(x)$ are the ultra-spherical Bessel functions [6] defined recursively from

$$\frac{d}{dx}\Phi_\ell^\nu(x) = \ell \coth x \Phi_\ell^\nu(x) - [\nu^2 + (\ell + 1)^2]^{1/2} \Phi_{\ell+1}^\nu(x), \quad (\text{A.13})$$

where

$$\Phi_0^\nu(x) = \frac{\sin \nu x}{\nu \sinh x}. \quad (\text{A.14})$$

We choose the normalisation constant $N(\nu)$ so that

$$\int d\Omega_{e^a|_R} dx \sinh^2 x Q_{a_1 a_2}^{(k)} Q^{(k') a_1 a_2} = \frac{\pi}{2} \delta_{\ell \ell'} \Delta_\ell \mathcal{Q}_{A_\ell} \mathcal{Q}'^{A_\ell} \nu^{-2} \delta(\nu - \nu'), \quad (\text{A.15})$$

which gives

$$N(\nu) = \frac{1}{\nu \sqrt{2(\nu^2 + 1)}} \left[\frac{(\ell + 2)!}{(\ell - 2)!} \right]^{1/2}. \quad (\text{A.16})$$

For closed universes, the hyperbolic functions should be replaced by their trigonometric counterparts in equations (A.9–A.16), $\nu^2 + n$ should be replaced by $\nu^2 - n$ with n an integer, and $\delta(\nu - \nu')$ should be replaced by $\delta_{\nu\nu'}$. For closed models, the regular, normalisable modes have ν an integer ≥ 3 , restricted to $\nu > \ell$.

Appendix A.2. Magnetic parity harmonics

For the magnetic parity tensor harmonics, which we denote with an overbar, we start with the general magnetic parity, rank-two PSTF tensor [47]:

$$\bar{Q}_{ab}^{(k)} = \bar{T}_1(x) [e_{c_\ell} \epsilon^{c_\ell c_{\ell-1}} ({}_a \mathcal{Q}_b)_{C_{\ell-1}} e^{C_{\ell-2}}]^{\text{TT}} + \bar{T}_2(x) e_d e_{(a} \epsilon_{b)}^{d c_\ell} \mathcal{Q}_{C_\ell} e^{C_{\ell-1}}, \quad (\text{A.17})$$

with $\ell \geq 2$. Equations (A.1) and (A.2) give

$$\bar{T}_1(x) = \frac{-1}{(\ell + 2) \sinh^2 x} \frac{d}{dx} [\sinh^3 x \bar{T}_2(x)], \quad (\text{A.18})$$

and

$$\frac{d^2}{dx^2} \bar{T}_2(x) + 4 \coth x \frac{d}{dx} \bar{T}_2(x) + \left[\nu^2 + 4 - \frac{(\ell^2 + \ell - 2)}{\sinh^2 x} \right] \bar{T}_2(x) = 0. \quad (\text{A.19})$$

The regular, normalisable solution of (A.19) is

$$\bar{T}_2(x) = \bar{N}(\nu) \frac{\Phi_\ell^\nu(x)}{\sinh x}, \quad (\text{A.20})$$

where $\nu \geq 0$. With the same normalisation as the electric parity harmonics, we find

$$\bar{N}(\nu) = \frac{1}{(\ell+1)} \left[\frac{2}{(\nu^2+1)} \right]^{1/2} \left[\frac{(\ell+2)!}{(\ell-2)!} \right]^{1/2}, \quad (\text{A.21})$$

so that $\bar{N}(\nu)/N(\nu) = 2\nu/(\ell+1)$. In a closed universe we make the same replacements as for the electric parity harmonics.

As a consequence of the first-order identity

$$D^2(\text{curl } S_{ab}) = \text{curl } (D^2 S_{ab}), \quad (\text{A.22})$$

where S_{ab} is an $O(\epsilon)$ PSTF tensor, it follows that if $Q_{ab}^{(k)}$ solves equation (A.1) then so does $\text{curl } Q_{ab}^{(k)}$. Furthermore, by virtue of equations (3.8) and (3.10), $\text{curl } Q_{ab}^{(k)}$ will also satisfy (A.2) if $Q_{ab}^{(k)}$ does. Since the curl operation is parity reversing, we expect the electric and magnetic parity tensor harmonics for given ℓ to be related by a curl. With the normalisation conventions adopted here, it is straightforward to show that

$$\text{curl } Q_{ab}^{(k)} = \frac{k}{S} \left(1 + \frac{3K}{k^2} \right)^{1/2} \bar{Q}_{ab}^{(k)}, \quad (\text{A.23})$$

$$\text{curl } \bar{Q}_{ab}^{(k)} = \frac{k}{S} \left(1 + \frac{3K}{k^2} \right)^{1/2} Q_{ab}^{(k)}, \quad (\text{A.24})$$

which are consistent with the first-order identity (3.13).

Appendix A.3. Sum over harmonic modes

A normalisable, first-order, rank-2 PSTF tensor field, such as the electric part of the Weyl tensor, can now be expanded in terms of the electric and magnetic parity tensor harmonics. Since the $\mathcal{Q}_{A_\ell}$ have only $2\ell+1$ degrees of freedom, it is convenient to introduce a set of $2\ell+1$ orthogonal basis PSTF tensors $\{\mathcal{Q}_{A_\ell}^{(\ell m)}\}$, with $m = -\ell, \dots, \ell$. We choose the basis tensors so that

$$\mathcal{Q}_{A_\ell}^{(\ell m)} \mathcal{Q}^{(\ell m') A_\ell} = \Delta_\ell^{-1} \delta_{mm'}, \quad (\text{A.25})$$

which implies

$$\sum_m \mathcal{Q}_{A_\ell}^{(\ell m)} \mathcal{Q}^{(\ell m) B_\ell} = \Delta_\ell^{-1} h_{\langle A_\ell}^{\langle B_\ell \rangle}. \quad (\text{A.26})$$

The tensor harmonic expansion of E_{ab} can then be written as

$$E_{ab} = \int_0^\infty |K|^{3/2} \nu^2 d\nu \sum_{\ell=2}^\infty \sum_{m=-\ell}^\ell (E_{\nu\ell m} Q_{ab}^{(k)} + \bar{E}_{\nu\ell m} \bar{Q}_{ab}^{(k)}), \quad (\text{A.27})$$

in an open universe. [Equation (A.27) can easily be generalised to include supercurvature modes, if these are present in the initial conditions.] In the text we use the abbreviated notation E_k for $E_{\nu\ell m}$, as we have done with the labelling of the tensor harmonics themselves. The summation and integral in (A.27) defines the symbolic summation over harmonic modes, $\sum_k$, employed in the text. In a closed universe, equation (A.27) should be replaced by

$$E_{ab} = \sum_{\nu=3}^\infty |K|^{3/2} \nu^2 \sum_{\ell=2}^{\nu-1} \sum_{m=-\ell}^\ell (E_{\nu\ell m} Q_{ab}^{(k)} + \bar{E}_{\nu\ell m} \bar{Q}_{ab}^{(k)}). \quad (\text{A.28})$$

Under the assumptions of statistical homogeneity and isotropy, the covariance structure of the E_k and $\bar{E}_k$ is restricted to the form

$$\begin{aligned}\langle E_k E_{k'} \rangle &= E^2(\nu) \delta_{kk'} \\ \langle \bar{E}_k \bar{E}_{k'} \rangle &= E^2(\nu) \delta_{kk'} \\ \langle E_k \bar{E}_{k'} \rangle &= 0,\end{aligned}\tag{A.29}$$

where the symbolic $\delta_{kk'}$ represents $\delta_{\ell\ell'}\delta_{mm'}|K|^{-3/2}\nu^{-2}\delta(\nu - \nu')$ in an open universe. In the closed case $\delta(\nu - \nu')$ should be replaced by $\delta_{\nu\nu'}$ in this expression.

References

- [1] Pen U, Seljak U and Turok N 1997 *Phys. Rev. Lett.* **79** 1611
- [2] Sachs R K and Wolfe A M 1967 *Ap. J.* **147** 73
- [3] Peebles P J E and Yu J T 1970 *Ap. J.* **162** 815
- [4] Wilson M L 1983 *Ap. J.* **273** L2
- [5] Bond J R and Efstathiou G 1984 *Ap. J.* **285** L45
- [6] Abbott L F and Schaefer R K 1986 *Ap. J.* **308** 546
- [7] Ma C P and Bertschinger E 1995 *Ap. J.* **455** 7
- [8] Seljak U and Zaldarriaga M 1996 *Ap. J.* **469** 437
- [9] Dautcourt G 1967 *Mon. Not. R. Astron. Soc.* **144** 255
- [10] Rubakov V A, Sazhin M V and Veyaskin A V 1982 *Phys. Lett.* **115B** 189
- [11] Fabbri R and Pollock M D 1983 *Phys. Lett.* **125B** 445
- [12] Abbott L F and Wise M B 1984 *Nucl. Phys.* **B244** 541
- [13] White M 1992 *Phys. Rev. D* **46** 4198
- [14] Crittenden R, Bond J R, Davis R L, Efstathiou G and Steinhardt P J 1993 *Phys. Rev. Lett.* **71** 324
- [15] Atrio-Barandela F and Silk J 1994 *Phys. Rev. D* **49** 1126
- [16] Frewin R A, Polnarev A G and Coles P 1994 *Mon. Not. R. Astron. Soc.* **266** L21
- [17] Allen B, Caldwell R and Koranda S 1995 *Phys. Rev. D* **51** 1553
- [18] Hu W and White M 1997 *Ap. J.* **46** L1
- [19] Durrer R and Kahniashvili T 1998 *Hel. Phys. Acta* **71** 445
- [20] Lifshitz E M 1946 *J. Phys. (Moscow)* **10** 116
- [21] Bardeen J M 1980 *Phys. Rev. D* **22** 1882
- [22] Hu W and White M 1997 *Phys. Rev. D* **56** 596
- [23] Hu W, Seljak U, White M and Zaldarriaga M 1998 *Phys. Rev. D* **57** 3290
- [24] Ellis G F R and Bruni M 1989 *Phys. Rev. D* **40** 1804
- [25] Bruni M and Sonego S 1999 *Class. Quantum Grav.* **16** L29
- [26] Ellis G F R, Hwang J and Bruni M 1989 *Phys. Rev. D* **40** 1819
- [27] Ehlers J 1993 *Gen. Rel. Grav.* **25** 1225
- [28] Maartens R, Ellis G F R and Stoeger W R 1995 *Phys. Rev. D* **51** 1525
- [29] Stoeger W R, Maartens R and Ellis G F R 1995 *Ap. J.* **443** 1
- [30] Dunsby P K S 1997 *Class. Quantum Grav.* **14** 3391
- [31] Challinor A D and Lasenby A N 1998 *Phys. Rev. D* **58** 023001
- [32] Challinor A D and Lasenby A N 1999 *Ap. J.* **513** 1
- [33] Gebbie T and Ellis G F R 1999 *Ann. Phys.* in press (astro-ph/9804316)
- [34] Maartens R, Gebbie T and Ellis G F R 1999 *Phys. Rev. D* **59** 083506
- [35] Challinor A D 1999 *Gen. Rel. Grav.* in press (astro-ph/9903283)
- [36] Gebbie T, Dunsby P K S and Ellis G F R 1999 *Ann. Phys.* in press (astro-ph/9904408)
- [37] Dunsby P K S, Bassett B A C C and Ellis G F R 1997 *Class. Quantum Grav.* **14** 1215

- [38] Ellis G F R and Elst H V 1999 *Theoretical and Observational Cosmology* ed M. Lachiéze-Rey (Dordrecht: Kluwer) p 1
- [39] Maartens R and Bassett B A 1998 *Class. Quantum Grav.* **15** 1998
- [40] Zakharov V D 1973 *Gravitational radiation in Einstein's theory* (New York: Halsted/Wiley) p 51
- [41] Challinor A D 1999 *Preprint* astro-ph/9911481
- [42] Hawking S W 1966 *Ap. J.* **145** 544
- [43] Albrecht A, Ferreira P, Joyce M and Prokopec T 1994 *Phys. Rev. D* **50** 4807
- [44] Polarski D and Starobinsky A A 1996 *Class. Quantum Grav.* **13** 377
- [45] Bucher M and Cohn J D 1997 *Phys. Rev. D* **55** 7461
- [46] García-Bellido J, Garriga J and Montes X 1999 *Phys. Rev. D* **60** 083501
- [47] Thorne K S 1980 *Rev. Mod. Phys.* **52** 299